	HEST MATION DISCLOSURE CITATION		
Attorney Docket No.: GC781-2-US	10 4/2	Serial No.: 10/530,556	
Applicant: Dunn-Coleman, et al.	JUL 0 2 2007 B		
Filing Date: November 5, 2003	300	Group:	
Page _ 1 _ of _ 5	TATE TO A	Date of this Submission: June 28, 2007	

Č.

US PATENT DOCUMENTS

Examiner's	Document				Sub-	Filing
Initial	Number	Date	Name	Class	Class	Date
	4,435,307	3/6/84	Barbesgaard			
	4,816,567	3/28/89	Cabilly			
	5,648,263	7/15/97	Schulein			
	5,691,178	11/25/97	Schulein			
	5,776,757	7/7/98	Schulein			
	6,022,725	2/8/00	Fowler			
	6,184,018	2/6/01	Li			
	6,982,159	9/21/01	Dunn-Coleman			
	7,045,332	12/18/01	Dunn-Coleman			
	7,005,289	12/18/01	Dunn-Coleman			1
	Patent Application No.: 20040102619		Dunn-Coleman			11/21/02

FOREIGN PATENT DOCUMENTS

Examiner's	Document				Sub-	Translation
Initials	Number	Date	Country	Class	Class	Yes/No
	WO 00/70031		PCT			
	WO 91/04673		PCT			
***************************************	WO 92/06209		PCT			
	WO 94/28117		PCT			
•	WO 96/02551	T	PCT			
	1,358,599		GB		-	
	2,094,826	3/2/82	GB			
	2,095,275	3/2/82	GB			

OTHER DOCUMENTS

Examiner's	Author, Title, Date, Pertinent Pages, etc.	
Initials		
	Altschul, S. F., et al., J. Mol. Biol. 215:403-410, 1990. Basic Local Alignment Search Tool	
	Altschul, S. F., et al., Nucleic Acids Res. 25:3389-3402, 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs.	

		I	
Attorney Docket No.: GC781-2-US		Serial No.: 10/530,556	
	unn-Coleman, et al.		
	lovember 5, 2003	Group:	
Page _2_ of	f5	Date of this Submission: June 28, 2007	
		s):24309-14. (Epub 2001 Apr 13.) ACELL, a novel ation of Cellulase and Xylanase Genes of Trichoderma	
	Aubert J.P. et al., Academic Press pp. 71- Utilization.	-86 (1988) Scale Up Of Cellulase Production and	
**	1993.	olecular Biology, John Wiley & Sons, New York, NY,	
	filter and microarray techniques to detect		
	Encoding An Extracelluar B-Glucosidase	62-567, 1991. Cloning and Amplification of the Gene From Trichoderma Reesei: Evidence for Improved	
**	Baulcombe, D., Arch. Virol. Suppl. 15:189	9-201, 1999.	
	Bennett, J.W. and Lasure, L.L. (eds.) Mon 396-428	e Gene Manipulations in Fungi. Academic Press, pp.	
	Campbell, et al., (1989) Curr. Genet. 16:5 Aspergillus niger using the homologous n	3-56, 1989. Improved transformation efficiency of iaD gene for nitrate reductase.	
	Chen et al., Biochem. Biophys. Acta. 1121:54-60, 1992. Purification and Characterization of two extracellular B-glucosidases from Trichoderma reesei		
••	Coligan, J. E. et al., eds., CURRENT PROTO	DCOLS IN IMMUNOLOGY, 1991.	
	Collen, A., et al., Journal of Chromatograp peptide fusions for improved protein partit	ohy A 910:275-284, 2001. Genetically engineered ioning in aqueous two-phase systems.	
**	Coughlan, et al., BIOCHEMISTRY AND GENE	TICS OF CELLULOSE DEGRADATION	
	Cummings and Fowler, Curr. Genet. 29:2 glucosidase by Saccharomyces cerevisia	27-233, 1996. Secretion of Trichoderma reesei B- e.	
	Dayhoff et al. in Atlas of Protein Sequence pp. 345-352, 1978. A Model Of Evolutional	e and Structure, Volume 5, Supplement 3, Chapter 22, ary Change in Proteins	
	Fields and Song, Nature 340:245-246, 19 Interactions.	89. A Novel Genetic System To Detect Protein-Protein	
	from solid-state cultures of Humicola grise		
	Enzymes in the Filamentous Fungus Trick		
	Freer, et al. J. Biol. Chem. 268:9337-9342 from a Yeast, Candida wickerhamii.	2, 1993. Kinetic Characterization of a B-Glucosidase	
**	Freshney, R. I., ed., ANIMAL CELL CULTURE, 1987.		
	Fusarium include Bajar, Podila and Kolattukudy, (1991) Proc. Natl. Acad. Sci. USA 88: 8202-8212		
	Goyal,et al. Bioresource Technol. 36-37-5	0 (1991). Characteristics of Fungal Cellulases.	
	overexpression of a Rhodothermus marin hydrolase family 12.	echnol. 49(3):277-84, 1998. Cloning, sequencing and us gene encoding a thermostable cellulose of glycosyl	
	of glycosyl hydrolases based on amino ac		
	Herr et al., Appl. Microbiol. Biotechnol. 5:2 Extracellular B-Glucosidase from Lenzites	29-36, 1978. Purification and Properties Of An s Trabea.	

Page 3 of 5 Date of this Submission: June 28, 2007 Hu et al., Mol Cell Biol. vol.11, no. 11, pp. 5792-5799, 1991. Antibodies Specific For the Human Retionblastoma Protein Identify A Family Of Related Polypeptides. Ilmen, M. et al., Appl. Environ. Microbiol. 63:1298-1306 (1997). Regulation of Cellulase Gene Expression in the Filamentous Fungus Trichoderma resest. Jakobovits, A, Curr Opin Biotechnol (5(5):561-6, 1995. Production of Fully Human Antibodies by Transgenic mice. Jakobovits, A, et al., Ann NY Acad Sci 764:525-35, 1995. Production of Antigen-Specific Human Antibodies tom Mice Engineered With Human Heavy and Light Chain YAC's. Jones et al., Nature 321:522-525, 1986. Replacing the complementarity determining regions in a human antibody with those from a mouse. Kawaguchi, T et al., Gene 173(2):287-8, 1996. Cloning and sequencing of the cDNA encoding B-glucosidase 1 from Aspergillus aculeatus. Kelley et al. ENBO J. 4 475-479, 1985. Transformation of Aspergillus niger by the amdS gene of Aspergillus nidulans. Knowles, J. et al., TiBTECH 5, 255-261, 1987. Cellulase families and their genes. Kohler and Milstein, Nature 26:495, 1975. Continuous cultures of fused cells secreting antibody of predefined specificity. Krishna, S. et al., Tiescource Tech. 77:193-196, 2001. Simultaneous saccharification and fermentation of lipnocellilulosic wastes to ethanol using a thermotolerant yeast. Kumar, A., et al., Textile Chemist and Colorist 29:37-42, 1997. Optimizing the use of cellulose enzymes in finishing cellulosic tabrics. Lehtio, J. et al., Textile Chemist and Colorist 29:37-42, 1997. Optimizing the use of cellulose enzymes in finishing cellulosic tabrics. Lehtio, J. et al., Textile Chemist and Colorist 29:37-42, 1997. Optimizing the use of cellulose enzymes in finishing cellulosic tabrics. Lehtio, J. et al., Textile Chemist and Colorist 29:37-42, 1997. Optimizing the use of cellulose-binding domain. Li and Ljungdahl Appl. Environ. Microbiol 62:209-213, 1996. Expression of Aureobasidium pullulans xynA	Applicant: Di	unn-Coleman, et al.	
Hu et al., Mol Cell Biol. vol.11, no. 11, pp. 5792-5799, 1991. Antibodies Specific For the Human Retionblastoma Protein Identify A Family Of Related Polypeptides. Ilmen, M. et al., Appl. Environ. Microbiol. 63:1298-1306 (1997). Regulation of Cellulase Gene Expression in the Filamentous Fungus Trichoderma ressei. Jakobovits, A. Curr Opin Biolechnol §(5):561-6, 1995. Production of Fully Human Antibodies by Transgenic mice. Jakobovits, A., et al., Ann NY Acad Sci 764:525-35, 1995. Production of Antigen-Specific Human Antibodies from Mice Engineered With Human Heavy and Light Chain YAC's. Jones et al., Nature 321:522-525, 1986. Replacing the complementarity determining regions in a human antibody with those from a mouse. Kawaguchi, T et al., Gene 173(2):287-8, 1996. Cloning and sequencing of the cDNA encoding B-glucosidase I from Aspergillus aculeatus. Kelley et al. EMBO J. 4:475-479, 1985. Transformation of Aspergillus niger by the amdS gene of Aspergillus indulans Knowles, J. et al., TIBTECH 5, 255-261, 1987. Cellulase families and their genes. Kohler and Milstein, Nature 256:495, 1975. Continuous cultures of fused cells secreting antibody of predefined specificity. Krishna, S. et al., Bioresource Tech. 77:193-196, 2001. Simultaneous saccharification and fermentation of lignocellulosic wastes to ethanol using a thermotolerant yeast. Kumar, A., et al., Textile Chemist and Colorist 29:37-42, 1997. Optimizing the use of cellulose enzymes in finishing cellulosic fabrics. Lehtio, J. et al., FEMS Microbiology Letters 195:197-204, 2001. Directed immobilization of recombinant staphylococci on cotton fibers by functional display of a fungal cellulose-binding domain. Li and Ljungdahl Appl. Environ. Microbiol. 62:209-213, 1996. Expression of Aureobasidium pullulans xynh in, and Secretion of the Xylanase from, Saccharomyces cerevisiae. Linder, M. and Teeri, T.T., Biotechnol. 57:15-28, 1997. Lorito, Hayes, DiPietro and Harman (1993) Curr. Genet. 24: 349-356; Medv. J. et al., Science, 249:915-918, 1990. A B3 Int	Filing Date: N	November 5, 2003	Group:
Retionblastoma Protein Identify A Family Of Related Polypeptides. Illmen, M. et al., Appl. Environ. Microbiol. 63:1298-1306 (1997). Regulation of Cellulase Gene Expression in the Filamentous Fungus Trichoderma reesel. Jakobovits, A, Curr Opin Biotechnol 6(5):561-6, 1995. Production of Fully Human Antibodies by Transgenic mice. Jakobovits, A, et al., Ann NY Acad Sci 764:525-35, 1995. Production of Antigen-Specific Human Antibodies from Mice Engineered With Human Heavy and Light Chain YAC. S. Jones et al., Nature 321:522-525, 1986. Replacing the complementarity determining regions in a human antibody with those from a mouse. Kawaguchi, T et al., Gene 173(2):287-8, 1996. Cloning and sequencing of the cDNA encoding B-glucosidase 1 from Aspergillus aculeatus. Kelley et al. EMBO J. 4:475-479, 1985. Transformation of Aspergillus niger by the amdS gene of Aspergillus nidulans Knowles, J. et al., TIBTECH 5, 255-261, 1987. Cellulase families and their genes. Kohler and Milstein, Nature 256:495, 1975. Continuous cultures of fused cells secreting antibody of predefined specificity. Krishna, S. et al., Bioresource Tech. 77:193-196, 2001. Simultaneous saccharification and fermentation of lignocelllulosic wastes to ethanol using a thermotolerant yeast. Kumar, A., et al., Textile Chemist and Colorist 29:37-42, 1997. Optimizing the use of cellulose enzymes in finishing cellulosic fabrics. Lehtio, J. et al., Fettile Chemist and Colorist 29:37-42, 1997. Optimizing the use of cellulose enzymes in finishing cellulosic rabrics. Lehtio, J. et al., Fettile Chemist and Colorist 29:37-42, 1997. Optimizing the use of recombinant staphylococci on cotton fibers by functional display of a fungal cellulose-binding domain. Li and Ljungdahl Appl. Environ. Microbiol. 62:209-213, 1996. Expression of Aureobasidium pullulans xynA in, and Secretion of the Kylanase from, Saccharomyces cerevisiae. Linder, M. and Teeri, T.T., Biotechnol. 57:15-28, 1997. Loftus, J. et al., J. Chromatography A 808:153, 1998. Ion-exchange chromatographic p	Page3 o	f_5_	Date of this Submission: June 28, 2007
Retionblastoma Protein Identify A Family Of Related Polypeptides. Illmen, M. et al., Appl. Environ. Microbiol. 63:1298-1306 (1997). Regulation of Cellulase Gene Expression in the Filamentous Fungus Trichoderma reesel. Jakobovits, A, Curr Opin Biotechnol 6(5):561-6, 1995. Production of Fully Human Antibodies by Transgenic mice. Jakobovits, A, et al., Ann NY Acad Sci 764:525-35, 1995. Production of Antigen-Specific Human Antibodies from Mice Engineered With Human Heavy and Light Chain YAC. S. Jones et al., Nature 321:522-525, 1986. Replacing the complementarity determining regions in a human antibody with those from a mouse. Kawaguchi, T et al., Gene 173(2):287-8, 1996. Cloning and sequencing of the cDNA encoding B-glucosidase 1 from Aspergillus aculeatus. Kelley et al. EMBO J. 4:475-479, 1985. Transformation of Aspergillus niger by the amdS gene of Aspergillus nidulans Knowles, J. et al., TIBTECH 5, 255-261, 1987. Cellulase families and their genes. Kohler and Milstein, Nature 256:495, 1975. Continuous cultures of fused cells secreting antibody of predefined specificity. Krishna, S. et al., Bioresource Tech. 77:193-196, 2001. Simultaneous saccharification and fermentation of lignocelllulosic wastes to ethanol using a thermotolerant yeast. Kumar, A., et al., Textile Chemist and Colorist 29:37-42, 1997. Optimizing the use of cellulose enzymes in finishing cellulosic fabrics. Lehtio, J. et al., Fettile Chemist and Colorist 29:37-42, 1997. Optimizing the use of cellulose enzymes in finishing cellulosic rabrics. Lehtio, J. et al., Fettile Chemist and Colorist 29:37-42, 1997. Optimizing the use of recombinant staphylococci on cotton fibers by functional display of a fungal cellulose-binding domain. Li and Ljungdahl Appl. Environ. Microbiol. 62:209-213, 1996. Expression of Aureobasidium pullulans xynA in, and Secretion of the Kylanase from, Saccharomyces cerevisiae. Linder, M. and Teeri, T.T., Biotechnol. 57:15-28, 1997. Loftus, J. et al., J. Chromatography A 808:153, 1998. Ion-exchange chromatographic p			
Expression in the Filamentous Fingus Trichoderma reesei. Jakobovits, A. cur Opin Biotechnol 6(5):561-6, 1995. Production of Fully Human Antibodies by Transgenic mice. Jakobovits, A., et al., Ann NY Acad Sci 764:525-35, 1995. Production of Antigen-Specific Human Antibodies from Mice Engineered With Human Heavy and Light Chain YAC's. Jones et al., Nature 321:522-525, 1986. Replacing the complementarity determining regions in a human antibody with those from a mouse. Kawaguchi, T et al., Gene 173(2):287-8, 1996. Cloning and sequencing of the cDNA encoding B-glucosidase 1 from Aspergillus aculeatus. Kelley et al. EMBO J. 4:475-479, 1985. Transformation of Aspergillus niger by the amdS gene of Aspergillus niculans Knowles, J. et al., TIBTECH 5, 255-261, 1987. Cellulase families and their genes. Kohler and Milstein, Nature 256:495, 1975. Continuous cultures of fused cells secreting antibody of predefined specificity. Krishna, S. et al., Bioresource Tech. 77:193-196, 2001. Simultaneous saccharification and fermentation of lignocelliulosic wastes to ethanol using a thermotolerant yeast. Kumar, A., et al., Textle Chemist and Colorist 29:37-42, 1997. Optimizing the use of cellulose enzymes in finishing cellulosic fabrics. Lehtio, J. et al., FEMS Microbiology Letters 195:197-204, 2001. Directed immobilization of recombinant staphylococci on cotton fibers by functional display of a fungal cellulose-binding domain. Li and Ljungdahl Appl. Environ. Microbiol. 62:209-213, 1996. Expression of Aureobasidium pullulans xynA in, and Secretion of the Xylanase from, Saccharomyces cerevisiae. Linder, M. and Teeri, T.T., Biotechnol. 57:15-28, 1997. Loftus, J. et al., J. Chromatography A 808:153, 1988. Innexchange chromatographic purification and quantitative analysis of trichoderma reesei cellulases celloborydrolase I, II and endoglucanase II by fast protein liquid chromatography. Ohmiya et al., Biotechnol. Gen. Engineer. Rev. 14:365-414, 1997. Okada, M. et al., Appl. Environ. Microbiol., 64:555-563, 1988. Molecular Char			
Transgenic mice. Jakobovits, A, et al., Ann NY Acad Sci 764:525-35, 1995. Production of Antigen-Specific Human Antibodies from Mice Engineered With Human Heavy and Light Chain YAC's. Jones et al., Nature 321:522-525, 1986. Replacing the complementarity determining regions in a human antibody with those from a mouse. Kawaguchi, T et al., Gene 173(2):287-8, 1996. Cloning and sequencing of the cDNA encoding B-glucosidase 1 from Aspergillus aculeatus. Kelley et al. EMBO J. 4:475-479, 1985. Transformation of Aspergillus niger by the amdS gene of Aspergillus nidulans Knowles, J. et al., TIBTECH 5, 255-261, 1987. Cellulase families and their genes. Kohler and Milstein, Nature 256:495, 1975. Continuous cultures of fused cells secreting antibody of predefined specificity. Krishna, S. et al., Bioresource Tech. 77:193-196, 2001. Simultaneous saccharification and fermentation of lignocelllulosic wastes to ethanol using a thermotolerant yeast. Kumar, A., et al., Textile Chemist and Colorist 29:37-42, 1997. Optimizing the use of cellulose enzymes in finishing cellulosic fabrics. Lehtio, J. et al., FEMS Microbiology Letters 195:197-204, 2001. Directed immobilization of recombinant staphylococci on cotton fibers by functional display of a fungal cellulose-binding domain. Li and Ljungdahl Appl. Environ. Microbiol. 62:209-213, 1996. Expression of Aureobasidium pullulans xynA in, and Secretion of the Xylanase from, Saccharomyces cerevisiae. Linder, M. and Teeri, T.T., Biotechnol. 57:15-28, 1997. Loftus, J. et al., Science, 249:915-918, 1990. A B3 Integrin Mutation Abolishes Ligand Binding and Alters Divalent Cation-Dependent Conformation. Lorito, Hayes, DiPietro and Harman (1993) Curr. Genet. 24: 349-356; Medve, J. et al., J. Chromatography A 808:153, 1998. Ion-exchange chromatographic purification and quantitative analysis of trichoderma reesei cellulases celluloses celluloses sendonication of the Gene Encoding a Low-Molecular-Mass Endoglucanase from Trichoderma reesei QM9414. Ooi et al., Nucleic Acid Res. 18:5884			
Antibodies from Mice Engineered With Human Heavy and Light Chain YAC's. Jones et al., Nature 321:522-525, 1986. Replacing the complementarity determining regions in a human antibody with those from a mouse. Kawaguchi, T et al., Gene 173(2):287-8, 1996. Cloning and sequencing of the cDNA encoding B-glucosidase 1 from Aspergillus aculeatus. Kelley et al. EMBO J. 4:475-479, 1985. Transformation of Aspergillus niger by the amdS gene of Aspergillus nidulans Knowles, J. et al., TIBTECH 5, 255-261, 1987. Cellulase families and their genes. Kohler and Milistein, Nature 256:495, 1975. Continuous cultures of fused cells secreting antibody of predefined specificity. Krishna, S. et al., Bioresource Tech. 77:193-196, 2001. Simultaneous saccharification and fermentation of lignocelliulosic wastes to ethanol using a thermotolerant yeast. Kumar, A., et al., Textile Chemist and Colorist 29:37-42, 1997. Optimizing the use of cellulose enzymes in finishing cellulosic fabrics. Lehtio, J. et al., FEMS Microbiology Letters 195:197-204, 2001. Directed immobilization of recombinant staphylococci on cotton fibers by functional display of a fungal cellulose-binding domain. Li and Ljungdahl Appl. Environ. Microbiol. 62:209-213, 1996. Expression of Aureobasidium pullulans xynA in, and Secretion of the Xylanase from, Saccharomyces cerevisiae. Linder, M. and Teeri, T.T., Biotechnol. 57:15-28, 1997. Loftus, J. et al., Science, 249:915-918, 1990. A B3 Integrin Mutation Abolishes Ligand Binding and Allers Divalent Catlon-Dependent Conformation. Lorito, Hayes, DiPietro and Harman (1993) Curr. Genet. 24: 349-356; Medve, J. et al., J. Chromatography A 808:153, 1998. Ion-exchange chromatographic purification and quantitative analysis of trichoderma reesei cellulases cellobiohydrolase I, Il and endoglucanase II by fast protein liquid chromatography. Ohmiya et al., Biotechnol. Gen. Engineer. Rev. 14:365-414, 1997. Okada, M. et al., Appl. Environ. Microbiol., 64:555-563, 1988. Molecular Characterization and Heterologous Expression of t			561-6, 1995. Production of Fully Human Antibodies by
human antibody with those from a mouse. Kawaguchi, T et al., Gene 173(2):287-8, 1996. Cloning and sequencing of the cDNA encoding B-glucosidase 1 from Aspergillus acqueatus. Kelley et al. EMBO J. 4:475-479, 1985. Transformation of Aspergillus niger by the amdS gene of Aspergillus nidulans Knowles, J. et al., TIBTECH 5, 255-261, 1987. Cellulase families and their genes. Kohler and Milstein, Nature 256:495, 1975. Continuous cultures of fused cells secreting antibody of predefined specificity. Krishna, S. et al., Bioresource Tech. 77:193-196, 2001. Simultaneous saccharification and fermentation of lignocelliulosic wastes to ethanol using a thermotolerant yeast. Kumar, A., et al., Textile Chemist and Colorist 29:37-42, 1997. Optimizing the use of cellulose enzymes in finishing cellulosic fabrics. Lehtio, J. et al., FEMS Microbiology Letters 195:197-204, 2001. Directed immobilization of recombinant staphylococci on cotton fibers by functional display of a fungal cellulose-binding domain. Li and Ljungdahl Appl. Environ. Microbiol. 62:209-213, 1996. Expression of Aureobasidium pullulans xynA in, and Secretion of the Xylanase from, Saccharomyces cerevisiae. Linder, M. and Teeri, T.T., Biotechnol. 57:15-28, 1997. Loftus, J. et al., Science, 249:915-918, 1990. A B3 Integrin Mutation Abolishes Ligand Binding and Alters Divalent Cation-Dependent Conformation. Lorito, Hayes, DiPietro and Harman (1993) Curr. Genet. 24: 349-356; Medve, J. et al., J. Chromatography A 808:153, 1998. Ion-exchange chromatographic purification and quantitative analysis of trichoderma reesei cellulases cellobiohydrolase I, Il and endoglucanase II by fast protein liquid chromatography. Ohmiya et al., Biotechnol. Gen. Engineer. Rev. 14:365-414, 1997. Okada, M. et al., Appl. Environ. Microbiol., 64:555-563, 1988. Molecular Characterization and Heterologous Expression of the Gene Encoding a Low-Molecular-Mass Endoglucanase from Trichoderma reesei QM9414. Ooi et al., Nucleic Acid Res. 18:5884, 1990. Complete nucleotide sequence of a gene		Antibodies from Mice Engineered With Hu	ıman Heavy and Light Chain YAC's.
glucosidase 1 from Aspergillus aculeatus. Kelley et al. EMBO J. 4:475-479, 1985. Transformation of Aspergillus niger by the amdS gene of Aspergillus nidulans Knowles, J. et al., TIBTECH 5, 255-261, 1987. Cellulase families and their genes. Kohler and Milstein, Nature 256:495, 1975. Continuous cultures of fused cells secreting antibody of predefined specificity. Krishna, S. et al., Bioresource Tech. 77:193-196, 2001. Simultaneous saccharification and fermentation of lignocelliulosic wastes to ethanol using a thermotolerant yeast. Kumar, A., et al., Textile Chemist and Colorist 29:37-42, 1997. Optimizing the use of cellulose enzymes in finishing cellulosic fabrics. Lehtio, J. et al., FEMS Microbiology Letters 195:197-204, 2001. Directed immobilization of recombinant staphylococci on cotton fibers by functional display of a fungal cellulose-binding domain. Li and Ljungdahl Appl. Environ. Microbiol. 62:209-213, 1996. Expression of Aureobasidium pullulans xynA in, and Secretion of the Xylanase from, Saccharomyces cerevisiae. Linder, M. and Teeri, T.T., Biotechnol. 57:15-28, 1997. Loftus, J. et al., Science, 249:915-918, 1990. A B3 Integrin Mutation Abolishes Ligand Binding and Alters Divalent Cation-Dependent Conformation. Lorito, Hayes, DiPietro and Harman (1993) Curr. Genet. 24: 349-356; Medve, J. et al., J. Chromatography A 808:153, 1998. Ion-exchange chromatographic purification and quantitative analysis of Inchoderma reesei cellulases cellobiohydrolase I, Il and endoglucanase II by fast protein liquid chromatography. Ohmity et al., Biotechnol. Gen. Engineer. Rev. 14:365-414, 1997. Okada, M. et al., Appl. Environ. Microbiol., 64:555-563, 1988. Molecular Characterization and Heterologous Expression of the Gene Encoding a Low-Molecular-Mass Endoglucanase from Trichoderma reesei CM9414. Ooi et al., Nucleic Acid Res. 18:5884, 1990. Complete nucleotide sequence of a gene coding for Aspergillus aculeatus cellulase (Fl-CMcase) Ortega et al., International Biodeterioration and Biodegradation 47:7-14, 2001		human antibody with those from a mouse.	
Aspergillus nidulans Knowles, J. et al., TIBTECH 5, 255-261, 1987. Cellulase families and their genes. Kohler and Milstein, Nature 256:495, 1975. Continuous cultures of fused cells secreting antibody of predefined specificity. Krishna, S. et al., Bioresource Tech. 77:193-196, 2001. Simultaneous saccharification and fermentation of lignocelliulosic wastes to ethanol using a thermotolerant yeast. Kumar, A., et al., Textile Chemist and Colorist 29:37-42, 1997. Optimizing the use of cellulose enzymes in finishing cellulosic fabrics. Lehtio, J. et al., FEMS Microbiology Letters 195:197-204, 2001. Directed immobilization of recombinant staphylococci on cotton fibers by functional display of a fungal cellulose-binding domain. Li and Ljungdahl Appl. Environ. Microbiol. 62:209-213, 1996. Expression of Aureobasidium pullulans xynA in, and Secretion of the Xylanase from, Saccharomyces cerevisiae. Linder, M. and Teeri, T.T., Biotechnol. 57:15-28, 1997. Loftus, J. et al., Science, 249:915-918, 1990. A B3 Integrin Mutation Abolishes Ligand Binding and Alters Divalent Catlon-Dependent Conformation. Lorito, Hayes, DiPietro and Harman (1993) Curr. Genet. 24: 349-356; Medve, J. et al., J. Chromatography A 808:153, 1998. Ion-exchange chromatographic purification and quantitative analysis of trichoderma reesei cellulases cellobiohydrolase I, II and endoglucanase II by fast protein liquid chromatography. Ohmiya et al., Biotechnol. Gen. Engineer. Rev. 14:365-414, 1997. Okada, M. et al., Appl. Environ. Microbiol., 64:555-563, 1988. Molecular Characterization and Heterologous Expression of the Gene Encoding a Low-Molecular-Mass Endoglucanase from Trichoderma reesei QM9414. Ooi et al., Nucleic Acid Res. 18:5884, 1990. Complete nucleotide sequence of a gene coding for Aspergillus aculeatus cellulase (FI-CMcase) Ortega et al., International Biodeterioration and Biodegradation 47:7-14, 2001. Kinetics of cellulose saccharification by Trichoderma reesei cellulases.	·	glucosidase 1 from Aspergillus aculeatus.	
Kohler and Milstein, Nature 256:495, 1975. Continuous cultures of fused cells secreting antibody of predefined specificity. Krishna, S. et al., Bioresource Tech. 77:193-196, 2001. Simultaneous saccharification and fermentation of lignocelliulosic wastes to ethanol using a thermotolerant yeast. Kumar, A., et al., Textile Chemist and Colorist 29:37-42, 1997. Optimizing the use of cellulose enzymes in finishing cellulosic fabrics. Lehtio, J. et al., FEMS Microbiology Letters 195:197-204, 2001. Directed immobilization of recombinant staphylococci on cotton fibers by functional display of a fungal cellulose-binding domain. Li and Ljungdahl Appl. Environ. Microbiol. 62:209-213, 1996. Expression of Aureobasidium pullulans xynA in, and Secretion of the Xylanase from, Saccharomyces cerevisiae. Linder, M. and Teeri, T.T., Biotechnol. 57:15-28, 1997. Loftus, J. et al., Science, 249:915-918, 1990. A B3 Integrin Mutation Abolishes Ligand Binding and Alters Divalent Cation-Dependent Conformation. Lorito, Hayes, DiPictro and Harman (1993) Curr. Genet. 24: 349-356; Medve, J. et al., J. Chromatography A 808:153, 1998. Ion-exchange chromatographic purification and quantitative analysis of trichoderma reesei cellulases cellobiohydrolase I, II and endoglucanase II by fast protein liquid chromatography. Ohmiya et al., Biotechnol. Gen. Engineer. Rev. 14:385-414, 1997. Okada, M. et al., Appl. Environ. Microbiol., 64:555-563, 1988. Molecular Characterization and Heterologous Expression of the Gene Encoding a Low-Molecular-Mass Endoglucanase from Trichoderma reesei QM9414. Ooi et al., Nucleic Acid Res. 18:5884, 1990. Complete nucleotide sequence of a gene coding for Aspergillus aculeatus cellulase (FI-CMcase) Ortega et al., International Biodeterioration and Biodegradation 47:7-14, 2001. Kinetics of cellulose saccharification by Trichoderma reesei cellulases.			ransformation of Aspergillus niger by the amdS gene of
of predefined specificity. Krishna, S. et al., Bioresource Tech. 77:193-196, 2001. Simultaneous saccharification and fermentation of lignocelliulosic wastes to ethanol using a thermotolerant yeast. Kumar, A., et al., Textile Chemist and Colorist 29:37-42, 1997. Optimizing the use of cellulose enzymes in finishing cellulosic fabrics. Lehtio, J. et al., FEMS Microbiology Letters 195:197-204, 2001. Directed immobilization of recombinant staphylococci on cotton fibers by functional display of a fungal cellulose-binding domain. Li and Ljungdahl Appl. Environ. Microbiol. 62:209-213, 1996. Expression of Aureobasidium pullulans xynA in, and Secretion of the Xylanase from, Saccharomyces cerevisiae. Linder, M. and Teeri, T.T., Biotechnol. 57:15-28, 1997. Loftus, J. et al., Science, 249:915-918, 1990. A B3 Integrin Mutation Abolishes Ligand Binding and Alters Divalent Cation-Dependent Conformation. Lorito, Hayes, DiPietro and Harman (1993) Curr. Genet. 24: 349-356; Medve, J. et al., J. Chromatography A 808:153, 1998. Ion-exchange chromatographic purification and quantitative analysis of trichoderma reesei cellulases cellobiohydrolase I, Il and endoglucanase II by fast protein liquid chromatography. Ohmiya et al., Biotechnol. Gen. Engineer. Rev. 14:385-414, 1997. Okada, M. et al., Appl. Environ. Microbiol., 64:555-563, 1988. Molecular Characterization and Heterologous Expression of the Gene Encoding a Low-Molecular-Mass Endoglucanase from Trichoderma reesei QM9414. Ooi et al., Nucleic Acid Res. 18:5884, 1990. Complete nucleotide sequence of a gene coding for Aspergillus aculeatus cellulase (FI-CMcase) Ortega et al., International Biodeterioration and Biodegradation 47:7-14, 2001. Kinetics of cellulose saccharification by Trichoderma reesei cellulase genes of Trichoderma reesei		<u> </u>	
fermentation of lignocelliulosic wastes to ethanol using a thermotolerant yeast. Kumar, A., et al., Textile Chemist and Colorist 29:37-42, 1997. Optimizing the use of cellulose enzymes in finishing cellulosic fabrics. Lehtio, J. et al., FEMS Microbiology Letters 195:197-204, 2001. Directed immobilization of recombinant staphylococci on cotton fibers by functional display of a fungal cellulose-binding domain. Li and Ljungdahl Appl. Environ. Microbiol. 62:209-213, 1996. Expression of Aureobasidium pullulans xynA in, and Secretion of the Xylanase from, Saccharomyces cerevisiae. Linder, M. and Teeri, T.T., Biotechnol. 57:15-28, 1997. Loftus, J. et al., Science, 249:915-918, 1990. A B3 Integrin Mutation Abolishes Ligand Binding and Alters Divalent Cation-Dependent Conformation. Lorito, Hayes, DiPietro and Harman (1993) Curr. Genet. 24: 349-356; Medve, J. et al., J. Chromatography A 808:153, 1998. Ion-exchange chromatographic purification and quantitative analysis of Inchoderma reesei cellulases cellobiohydrolase I, Il and endoglucanase II by fast protein liquid chromatography. Ohmity et al., Biotechnol. Gen. Engineer. Rev. 14:365-414, 1997. Okada, M. et al., Appl. Environ. Microbiol., 64:555-563, 1988. Molecular Characterization and Heterologous Expression of the Gene Encoding a Low-Molecular-Mass Endoglucanase from Trichoderma reesei QM9414. Ooi et al., Nucleic Acid Res. 18:5884, 1990. Complete nucleotide sequence of a gene coding for Aspergillus aculeatus cellulase (Fl-CMcase) Ortega et al., International Biodeterioration and Biodegradation 47:7-14, 2001. Kinetics of cellulose saccharification by Trichoderma reesei cellulases. Penttill et al., Gene 45:253-263, 1986. Homology between cellulase genes of Trichoderma reesei.		of predefined specificity.	
enzymes in finishing cellulosic fabrics. Lehtio, J. et al., FEMS Microbiology Letters 195:197-204, 2001. Directed immobilization of recombinant staphylococci on cotton fibers by functional display of a fungal cellulose-binding domain. Li and Ljungdahl Appl. Environ. Microbiol. 62:209-213, 1996. Expression of Aureobasidium pullulans xynA in, and Secretion of the Xylanase from, Saccharomyces cerevisiae. Linder, M. and Teeri, T.T., Biotechnol. 57:15-28, 1997. Loftus, J. et al., Science, 249:915-918, 1990. A B3 Integrin Mutation Abolishes Ligand Binding and Alters Divalent Catlon-Dependent Conformation. Lorito, Hayes, DiPietro and Harman (1993) Curr. Genet. 24: 349-356; Medve, J. et al., J. Chromatography A 808:153, 1998. Ion-exchange chromatographic purification and quantitative analysis of trichoderma reesei cellulases cellobiohydrolase I, II and endoglucanase II by fast protein liquid chromatography. Ohmiya et al., Biotechnol. Gen. Engineer. Rev. 14:365-414, 1997. Okada, M. et al., Appl. Environ. Microbiol., 64:555-563, 1988. Molecular Characterization and Heterologous Expression of the Gene Encoding a Low-Molecular-Mass Endoglucanase from Trichoderma reesei QM9414. Ooi et al., Nucleic Acid Res. 18:5884, 1990. Complete nucleotide sequence of a gene coding for Aspergillus aculeatus cellulase (FI-CMcase) Ortega et al., International Biodeterioration and Biodegradation 47:7-14, 2001. Kinetics of cellulose saccharification by Trichoderma reesei cellulases. Penttila et al., Gene 45:253-263, 1986. Homology between cellulase genes of Trichoderma reesei.		fermentation of lignocelllulosic wastes to e	ethanol using a thermotolerant yeast.
recombinant staphylococci on cotton fibers by functional display of a fungal cellulose-binding domain. Li and Ljungdahl Appl. Environ. Microbiol. 62:209-213, 1996. Expression of Aureobasidium pullulans xynA in, and Secretion of the Xylanase from, Saccharomyces cerevisiae. Linder, M. and Teeri, T.T., Biotechnol. 57:15-28, 1997. Loftus, J. et al., Science, 249:915-918, 1990. A B3 Integrin Mutation Abolishes Ligand Binding and Alters Divalent Cation-Dependent Conformation. Lorito, Hayes, DiPietro and Harman (1993) Curr. Genet. 24: 349-356; Medve, J. et al., J. Chromatography A 808:153, 1998. Ion-exchange chromatographic purification and quantitative analysis of Inchoderma reesei cellulases cellobiohydrolase I, II and endoglucanase II by fast protein liquid chromatography. Ohmiya et al., Biotechnol. Gen. Engineer. Rev. 14:385-414, 1997. Okada, M. et al., Appl. Environ. Microbiol., 64:555-563, 1988. Molecular Characterization and Heterologous Expression of the Gene Encoding a Low-Molecular-Mass Endoglucanase from Trichoderma reesei QM9414. Ooi et al., Nucleic Acid Res. 18:5884, 1990. Complete nucleotide sequence of a gene coding for Aspergillus aculeatus cellulase (FI-CMcase) Ortega et al., International Biodeterioration and Biodegradation 47:7-14, 2001. Kinetics of cellulose saccharification by Trichoderma reesei cellulase genes of Trichoderma reesei.		enzymes in finishing cellulosic fabrics.	
pullulans xynA in, and Secretion of the Xylanase from, Saccharomyces cerevisiae. Linder, M. and Teeri, T.T., Biotechnol. 57:15-28, 1997. Loftus, J. et al., Science, 249:915-918, 1990. A B3 Integrin Mutation Abolishes Ligand Binding and Alters Divalent Cation-Dependent Conformation. Lorito, Hayes, DiPictor and Harman (1993) Curr. Genet. 24: 349-356; Medve, J. et al., J. Chromatography A 808:153, 1998. Ion-exchange chromatographic purification and quantitative analysis of trichoderma reesei cellulases cellobiohydrolase I, Il and endoglucanase II by fast protein liquid chromatography. Ohmiya et al., Biotechnol. Gen. Engineer. Rev. 14:385-414, 1997. Okada, M. et al., Appl. Environ. Microbiol., 64:555-563, 1988. Molecular Characterization and Heterologous Expression of the Gene Encoding a Low-Molecular-Mass Endoglucanase from Trichoderma reesei QM9414. Ooi et al., Nucleic Acid Res. 18:5884, 1990. Complete nucleotide sequence of a gene coding for Aspergillus aculeatus cellulase (FI-CMcase) Ortega et al., International Biodeterioration and Biodegradation 47:7-14, 2001. Kinetics of cellulose saccharification by Trichoderma reesei cellulases. Pentitia et al., Gene 45:253-263, 1986. Homology between cellulase genes of Trichoderma reesei		recombinant staphylococci on cotton fiber	
Loftus, J. et al., Science, 249:915-918, 1990. A B3 Integrin Mutation Abolishes Ligand Binding and Alters Divalent Cation-Dependent Conformation. Lorito, Hayes, DiPietro and Harman (1993) Curr. Genet. 24: 349-356; Medve, J. et al., J. Chromatography A 808:153, 1998. Ion-exchange chromatographic purification and quantitative analysis of trichoderma reesei cellulases cellobiohydrolase I, II and endoglucanase II by fast protein liquid chromatography. Ohmiya et al., Biotechnol. Gen. Engineer. Rev. 14:365-414, 1997. Okada, M. et al., Appl. Environ. Microbiol., 64:555-563, 1988. Molecular Characterization and Heterologous Expression of the Gene Encoding a Low-Molecular-Mass Endoglucanase from Trichoderma reesei QM9414. Ooi et al., Nucleic Acid Res. 18:5884, 1990. Complete nucleotide sequence of a gene coding for Aspergillus aculeatus cellulase (FI-CMcase) Ortega et al., International Biodeterioration and Biodegradation 47:7-14, 2001. Kinetics of cellulose saccharification by Trichoderma reesei cellulases. Penttila et al., Gene 45:253-263, 1986. Homology between cellulase genes of Trichoderma reesei		pullulans xynA in, and Secretion of the Xy	lanase from, Saccharomyces cerevisiae.
and Alters Divalent Cation-Dependent Conformation. Lorito, Hayes, DiPietro and Harman (1993) Curr. Genet. 24: 349-356; Medve, J. et al., J. Chromatography A 808:153, 1998. Ion-exchange chromatographic purification and quantitative analysis of trichoderma reesei cellulases cellobiohydrolase I, II and endoglucanase II by fast protein liquid chromatography. Ohmiya et al., Biotechnol. Gen. Engineer. Rev. 14:365-414, 1997. Okada, M. et al., Appl. Environ. Microbiol., 64:555-563, 1988. Molecular Characterization and Heterologous Expression of the Gene Encoding a Low-Molecular-Mass Endoglucanase from Trichoderma reesei QM9414. Ooi et al., Nucleic Acid Res. 18:5884, 1990. Complete nucleotide sequence of a gene coding for Aspergillus aculeatus cellulase (FI-OMcase) Ortega et al., International Biodeterioration and Biodegradation 47:7-14, 2001. Kinetics of cellulose saccharification by Trichoderma reesei cellulases. Pentitia et al., Gene 45:253-263, 1986. Homology between cellulase genes of Trichoderma reesei.		Linder, M. and Teeri, T.T., Biotechnol. 57:	15-28, 1997.
Medve, J. et al., J. Chromatography A 808:153, 1998. Ion-exchange chromatographic purification and quantitative analysis of trichoderma reesei cellulases cellobiohydrolase I, II and endoglucanase II by fast protein liquid chromatography. Ohmiya et al., Biotechnol. Gen. Engineer. Rev. 14:365-414, 1997. Okada, M. et al., Appl. Environ. Microbiol., 64:555-563, 1988. Molecular Characterization and Heterologous Expression of the Gene Encoding a Low-Molecular-Mass Endoglucanase from Trichoderma reesei QM9414. Ooi et al., Nucleic Acid Res. 18:5884, 1990. Complete nucleotide sequence of a gene coding for Aspergillus aculealus cellulase (FI-CMcase) Ortega et al., International Biodeterioration and Biodegradation 47:7-14, 2001. Kinetics of cellulose saccharification by Trichoderma reesei cellulases. Pentilia et al., Gene 45:253-263, 1986. Homology between cellulase genes of Trichoderma reesei			
and quantitative analysis of trichoderma reesei cellulases cellobiohydrolase I, il and endoglucanase II by fast protein liquid chromatography. Ohmiya et al., Biotechnol. Gen. Engineer. Rev. 14:365-414, 1997. Okada, M. et al., Appl. Environ. Microbiol., 64:555-563, 1988. Molecular Characterization and Heterologous Expression of the Gene Encoding a Low-Molecular-Mass Endoglucanase from Trichoderma reesei QM9414. Ooi et al., Nucleic Acid Res. 18:5884, 1990. Complete nucleotide sequence of a gene coding for Aspergillus aculealus cellulase (FI-CMcase) Ortega et al., International Biodeterioration and Biodegradation 47:7-14, 2001. Kinetics of cellulose saccharification by Trichoderma reesei cellulases. Pentilia et al., Gene 45:253-263, 1986. Homology between cellulase genes of Trichoderma reesei.		Lorito, Hayes, DiPietro and Harman (1993	3) Curr. Genet. 24: 349-356;
Okada, M. et al., Appl. Environ. Microbiol., 64:555-563, 1988. Molecular Characterization and Heterologous Expression of the Gene Encoding a Low-Molecular-Mass Endoglucanase from Trichoderma reesel QM9414. Ooi et al., Nucleic Acid Res. 18:5884, 1990. Complete nucleotide sequence of a gene coding for Aspergillus aculealus cellulase (FI-CMcase) Ortega et al., International Biodeterioration and Biodegradation 47:7-14, 2001. Kinetics of cellulose saccharification by Trichoderma reesei cellulases. Penttila et al., Gene 45:253-263, 1986. Homology between cellulase genes of Trichoderma reesei	1	and quantitative analysis of trichoderma re	eesei cellulases cellobiohydrolase I, II and
Heterologous Expression of the Gene Encoding a Low-Molecular-Mass Endoglucanase from Trichoderma reesei QM9414. Ooi et al., Nucleic Acid Res. 18:5884, 1990. Complete nucleotide sequence of a gene coding for Aspergillus aculealus cellulase (FI-CMcase) Ortega et al., International Biodeterioration and Biodegradation 47:7-14, 2001. Kinetics of cellulose saccharification by Trichoderma reesei cellulases. Penttila et al., Gene 45:253-263, 1986. Homology between cellulase genes of Trichoderma reesei.		Ohmiya et al., Biotechnol. Gen. Engineer.	Rev. 14:365-414, 1997.
Aspergillus aculeatus cellulase (FI-CMcase) Ortega et al., International Biodeterioration and Biodegradation 47:7-14, 2001. Kinetics of cellulose saccharification by Trichoderma reesei cellulases. Penttila et al., Gene 45:253-263, 1986. Homology between cellulase genes of Trichoderma reesei		Heterologous Expression of the Gene End	
Ortega et al., International Biodeterioration and Biodegradation 47:7-14, 2001. Kinetics of cellulose saccharification by Trichoderma reesei cellulases. Penttila et al., Gene 45:253-263, 1986. Homology between cellulase genes of Trichoderma reesei			
Penttila et al., Gene 45:253-263, 1986. Homology between cellulase genes of Trichoderma reesei	*	Ortega et al., International Biodeterioration	n and Biodegradation 47:7-14, 2001. Kinetics of
		Penttila et al., Gene 45:253-263, 1986. H	
			fficient secretion of two fungal cellobiohydrolases by

ALL REFERENCES CONSIDERED EXCEPT WHERE LINED THROUGH. /TS/

Serial No.: 10/530,556

Attorney Docket No.: GC781-2-US

Attorney Docket No.: GC/81-2-US Serial No.: 10/530,556		
Applicant: Du	nn-Coleman, et al.	
Filing Date: November 5, 2003 Group:		
Page 4 of		Date of this Submission: June 28, 2007
	•	
	Saccharomyces cerevisiae.	
		pression of Two Trichoderma reesei Endoglucanase in
	the Yeast Saccharomyces cerevisiae.	, and the second
	Pere, J., et al., In Proc. Tappi Pulping Con	f., Nashville, TN, 27-31, pp. 693-696, 1996.
**	Pourquie, J. et al., BIOCHEMISTRY AND GEN	IETICS OF CELLULOSE DEGRADATION, eds.
	Riechmann et al., Nature 332:323-327, 19	88. Reshaping human antibodies for therapy.
	Saccharomyces ceresvisiae	Synthesis and Secretion of wheat a-amylase in
	Saarilahti et al., Gene 90:9-14, 1990. Cele carotovora subsp. Carotovara.	S: a novel endoglucanase identified from Erwinia
	Sakamoto et al., Curr. Genet. 27:435-439,	1995
	Saloheimo M, et al., Gene 63:11-22, 1988 reesei; the characterization of both gene a	. EGIII, a new endoglucanese from Trichoderma and enzyme.
	Saloheimo, A. et al., Molecular Microbiolog gene, eg15, from Trichoderma reesei isola	gy, 13:219-228, 1994. A novel, small endoglucanase lated by expression in yeast.
-		9:584-591, 1997. cDNA cloning of a Trichoderma oglucanase activity by expression in yeast.
	Schulein, Methods Enzymol., 160, 25, pag	es 234 et seq, 1988. Cellulase of Trichoderma reesei.
	Scopes, Methods Enzymol. 90: 479-91, 1 muscle extract.	982. Purification of all glycolytic enzymes from one
	Shoemaker, S. et al., Bio/Technology, 1:69 Cellobiohydrolase I Dervived from Trichod	91-696, 1983. Molecular cloning of Exo- lerma Reesei Strain I.27.
		923-30, 1994. Cloning and sequencing of a ng for a thermostable B-glucanase and its expression in
	Stahlberg, J. et al., Bio/Technol. 9:286-290 cellulose based on the two-domain structu	D, 1991. A New model for enzymatic hydrolysis of tre of cellobiohydrolase I.
**	Strathern <i>et al.</i> , eds. (1981) The Molecular Harbor Press, Plainview. N.Y.	Biology of the Yeast Saccharomyces, Cold Spring
	domains in the hydrolysis and modification	
	Teeri, T. et al., Gene, 51:43-52, 1987. Hornenzymes	nologous domains in Trichoderma reesei cellulytic
	xynB from the thermophilic	190:13-19, 2000. Codon optimization of xylanase gene
	The Asexual Spores of A. nidulans.	anization of a Gene Cluster Expressed Specifically in
	trichoderma reesei QM 9414.	5-581, 1988. Studies of the cellulolytic system of
	Tormo, J. et al., EMBO J. 15:5739-5751, 1 binding domain; a general mechanism for	996. Crystal structure of a bacterial family-III cellulose- attachment to cellulose.
		t 24:23-26, 1992. Improving the softness and surface
	appearance of cotton fabrics and garment. Van Repsburg et al. Yeast 14:67-76, 1998	B. Engineering Yeast for Efficent Cellulose Degradation
	van Nonoburg of al., 16ast 14.07-70, 1330	5. Engineering reastroi Emcent Centrose Degradation

ALL REFERENCES CONSIDERED EXCEPT WHERE LINED THROUGH. /TS/

Attorney Doo	ket No.: GC781-2-US	Serial No.: 10/530,556
Applicant: D	unn-Coleman, et al.	
Filing Date:	November 5, 2003	Group:
Page5c	of <u>5</u>	Date of this Submission: June 28, 2007
	Van Tilbeurgh, H. et al., FEBS I cellobiohydrolase I from Trichoc	Lett. 204:223-227, 1986. Limited proteolysis of the derma reesei.
	VanMontagu and Herrera-Estre	ella (1990) Curr. Genet. 17:169-174
	antilysozyme Activity.	534-1536, 1988. Reshaping Human Antibodies : Grafting an
	Warrington, et al., Genomics 13	803-808, 1992
	Wells et al., Gene 34:315, 1985 multiple mutations at defined sit	. Cassette mutagenesis; an efficient method for generation of es.
	Wells et al., Philos. Trans. R. Se formation in stabilization the tra	oc. London SerA 317:415, 1986. Importance of hydrogen-bond nsition state of subtilisin.
•	Wood et al., METHODS IN ENZYM	DLOGY
	Wood, Biochem. Soc. Trans., 1	3, pp. 407-410, 1985. Properties of cellulolytic enzyme systems.
Examiner	/Tekchand Saidha/	Date Considered 07/17/2008
	nitial if reference considered, whether o	r not citation is in conformance with MPEP 609; draw line through citation